

**Problem:** Winter Grain Mites (WGM) - *Penthaleus major*



**Hosts:** Kentucky bluegrass, perennial ryegrass, tall fescue and Chewings red fescue

**Description:** The young mites are reddish-orange but become black with reddish legs in later stages. There is also a dorsal anus surrounded by a reddish-orange spot. This turf pest is unique in that it feeds during the winter to early spring and spends the summer in the egg stage. Damage often appears from January through mid-March and is evidenced by a silvery, scorched appearance to the turf. The injury is often masked until the turf greens up in the spring. Mites feed on cloudy days or at night. On sunny days, WGM's can be found on the crown of grass plants or in the thatch layer.

WGM's feed by rasping the leaf surface and sucking up the plant sap released. The silvery effect is caused by the loss of chlorophyll and desiccation. WGM's are not harmed by short periods of sleet, ice, or frozen ground. They actively feed under snow.

**Recommendations:** Presence of this mite does not necessarily warrant spraying. However, if damage causes the turf to become unsightly, Cyhalothrin in liquid formulation (Scimitar, Spectracide Triazicide) or wettable powder formulation (Battle) are both labeled against mites in general.

**References:**

1. Destructive Turf Insects, Harry Niemczyk, HDN Books
2. Turfgrass Insects of the United States and Canada, Cornell University Press, pg 26-29
3. [Winter Grain Mites](#), K-State Research and Extension, Entomology Publication MF-2073

**Last Update:** 11/17/2023

---

Brand names appearing in this publication are for product identification purposes only. No endorsement is intended, nor is criticism implied of similar products not mentioned.

*“Knowledge for Life”*

Kansas State University Agricultural Experiment Station and Cooperative Extension Service